

#### PRODUCT INFORMATION

## HindIII

**#ER0501** 5000 u

**Lot:** Expiry Date:

5'...**A↓A G C T T**... 3'

3'... **T T C G A**↑**A** ...5'

Concentration: 10 u/µl

Source: Haemophilus influenzae Rd

Supplied with: 1 ml of 10X Buffer R

1 ml of 10X Buffer Tango

Store at -20°C

 $\mathbf{R} \ ] \ [37^{\circ}$ 

95%









In total 3 vials. BSA included

www.thermoscientific.com/fermentas

#### **RECOMMENDATIONS**

**1X Buffer R** (for 100% HindIII digestion)

10 mM Tris-HCl (pH 8.5), 10 mM MgCl<sub>2</sub>, 100 mM KCl, 0.1 mg/ml BSA.

#### Incubate at 37°C.

#### **Unit Definition**

One unit is defined as the amount of HindIII required to digest 1  $\mu$ g lambda DNA in 1 hour at 37°C in 50  $\mu$ I of recommended reaction buffer.

#### **Dilution**

Dilute with Dilution Buffer (#B19): 10 mM Tris-HCl (pH 7.4 at 25°C), 100 mM KCl, 1 mM EDTA, 1 mM DTT, 0.2 mg/ml BSA and 50% glycerol.

## **Double Digests**

Thermo Scientific Tango Buffer is provided to simplify buffer selection for double digests. 98% of Thermo Scientific restriction enzymes are active in a 1X or 2X concentration of Tango $^{\text{\tiny TM}}$  Buffer. Please refer to to

<u>www.fermentas.com/doubledigest</u> to choose the best buffer for your experiments.

1X Tango Buffer: 33 mM Tris-acetate (pH 7.9 at 37°C), 10 mM magnesium acetate, 66 mM potassium acetate, 0.1 mg/ml BSA.

## **Storage Buffer**

HindIII is supplied in: 10 mM Tris-HCl (pH 7.5 at 25°C), 250 mM KCl, 1 mM DTT, 0.1 mM EDTA, 0.2 mg/ml BSA and 50% glycerol.

## **Recommended Protocol for Digestion**

• Add:

nuclease-free water	16 µl
10X Buffer R	2 μΙ
DNA (0.5-1 μg/μl)	1 µl
Hindll	0.5-2 µl*

- Mix gently and spin down for a few seconds.
- Incubate at 37°C for 1-16 hours.

The digestion reaction may be scaled either up or down.

# **Recommended Protocol for Digestion of PCR Products Directly after Amplification**

• Add:

PCR reaction mixture 10  $\mu$ l (~0.1-0.5  $\mu$ g of DNA) nuclease-free water 18  $\mu$ l 10X Buffer R 2  $\mu$ l HindIII 1-2  $\mu$ l\*

- Mix gently and spin down for a few seconds.
- Incubate at 37°C for 1-16 hours.

#### **Thermal Inactivation**

HindIII is inactivated by incubation at 80°C for 20 min.

#### **ENZYME PROPERTIES**

#### **Enzyme Activity in Thermo Scientific REase Buffers, %**

В	G	0	R	Tango	2X Tango
0-20	20-50	0-20	100	50-100	50-100

#### **Methylation Effects on Digestion**

Dam: never overlaps — no effect. Dcm: never overlaps — no effect. CpG: never overlaps — no effect. EcoKI: never overlaps — no effect.

EcoBl: may overlap – cleavage impaired.

## **Stability during Prolonged Incubation**

A minimum of 0.1 units of the enzyme is required for complete digestion of 1  $\mu$ g of lambda DNA in 16 hours at 37°C.

## **Digestion of Agarose-embedded DNA**

A minimum 5 units of the enzyme is required for complete digestion of 1  $\mu g$  of agarose-embedded lambda DNA in 16 hours.

## **Number of Recognition Sites in DNA**

λ	ФХ174	pBR322	pUC57	pUC18/19	pTZ19R/U	M13mp18/19
6	0	1	1	1	1	1

For **CERTIFICATE OF ANALYSIS** see back page

<sup>\*</sup> This volume of the enzyme is recommended for preparations of standard concentrations (10 u/μl), whereas HC enzymes (50 u/μl) should be diluted with Dilution Buffer to obtain 10u/μl concentration.

#### **CERTIFICATE OF ANALYSIS**

## **Overdigestion Assay**

No detectable change in the specific fragmentation pattern is observed after a 160-fold overdigestion with HindIII (10 u/µg lambda DNA x 16 hours).

#### **Ligation/Recutting Assay**

After a 50-fold overdigestion (3 u/µg DNA x 17 hours) with Hindlll, more than 95% of the digested DNA fragments can be ligated at a 5'-termini concentration of 0.07 µM. More than 95% of these sites can be recut.

## **Labeled Oligonucleotide (LO) Assay**

No detectable degradation of single-stranded or doublestranded labeled oligonucleotides occurred during incubation with 10 units of Hindll for 4 hours.

#### **Blue/White Cloning Assay**

pUC57 was incubated with 10 units of HindIII for 16 hours. After religation and transformation, the background level of white colonies was <1%.

#### **Quality authorized by:**



Jurgita Zilinskiene

#### PRODUCT USE LIMITATION

This product is developed, designed and sold exclusively for research purposes and in vitro use only. The product was not tested for use in diagnostics or for drug development, nor is it suitable for administration to humans or animals.

Please refer to www.thermoscientific.com/fermentas for Material Safety Data Sheet of the product.

© 2011 Thermo Fisher Scientific Inc. All rights reserved. All trademarks are the property of Thermo Fisher Scientific Inc. and its subsidiaries.