


Anti-Human CD279 (PD-1) PE

Catalog Number: 12-2799

RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of unstimulated (blue histogram) or 3-day PHA-stimulated normal human peripheral blood cells (purple histogram) with Anti-Human CD279 (PD-1) PE. Viable cells, as determined by Fixable Viability Dye eFluor[®] 660, in the lymphocyte gate were used for analysis.

Product Information

Contents: Anti-Human CD279 (PD-1) PE

Catalog Number: 12-2799

Clone: eBioJ105 (J105)

Concentration: 5 μ L (0.5 μ g)/test

Host/Isotype: Mouse IgG1, kappa

Formulation: aqueous buffer, 0.09% sodium

azide, may contain carrier protein/stabilizer

Temperature Limitation: Store at 2-8°C. Do not freeze. Light sensitive material.

Batch Code: Refer to vial

Use By: Refer to vial

Caution, contains Azide

Description

The J105 monoclonal antibody reacts with the human PD-1 (programmed death-1), a 55 kDa member of the CD28 immunoglobulin superfamily. PD-1 contains the immunoreceptor tyrosine-based inhibitory motif (ITIM) and plays a key role in peripheral tolerance and autoimmune disease. PD-1 is expressed predominantly on activated T and B lymphocytes. Two novel members of the B7 family have been identified as the PD-1 ligands, PD-L1 (B7-H1) and PD-L2 (B7-DC). Evidence reported to date suggests overlapping functions for these two PD-1 ligands and their constitutive expression on some normal tissues and upregulation on activated antigen-presenting cells.

Costaining experiments suggest that eBioJ105 recognizes a different epitope than MIH4 (cat. 11-9969).

Applications Reported

This eBioJ105 (J105) antibody has been reported for use in flow cytometric analysis.

Applications Tested

This eBioJ105 (J105) antibody has been pre-titrated and tested by flow cytometric analysis of PHA stimulated human peripheral blood cells. This can be used at 5 μ L (0.5 μ g) per test. A test is defined as the amount (μ g) of antibody that will stain a cell sample in a final volume of 100 μ L. Cell number should be determined empirically but can range from 10^5 to 10^8 cells/test.

References

Iwai Y, Okazaki T, Nishimura H, Kawasaki A, Yagita H, Honjo T. Microanatomical localization of PD-1 in human tonsils. *Immunol Lett.* 2002 Oct 1;83(3):215-20. (**J105**, FC, PubMed)

Related Products

12-4714 Mouse IgG1 K Isotype Control PE (P3.6.2.8.1)

46-0047 Anti-Human CD4 PerCP-eFluor[®] 710 (SK3 (SK-3))

65-0864 Fixable Viability Dye eFluor[®] 660

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.ebioscience.com •
info@ebioscience.com