


Anti-Mouse CD130 APC

Catalog Number: 17-1302

Also known as: gp130, IL-6 Receptor beta, IL-6RB

RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of C57Bl/6 splenocytes with Anti-Mouse CD3e FITC (cat. 11-0031) and 0.125 ug of Rat IgG2a kappa Isotype Control APC (cat. 17-4321) (left) or 0.125 ug of Anti-Mouse CD130 APC (right). Cells in the lymphocyte gate were used for analysis.

Product Information



Contents: Anti-Mouse CD130 APC

Catalog Number: 17-1302

Clone: KGP130

Concentration: 0.2 mg/mL

Host/Isotype: Rat IgG2a, kappa



Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

Temperature Limitation: Store at 2-8°C. Do not freeze. Light-sensitive material.

Batch Code: Refer to vial

Use By: Refer to vial

Description

The KGP130 monoclonal antibody reacts with mouse gp130, also known as CD130. gp130 is a 130 kD Type-I transmembrane glycoprotein containing a 597 amino acid extracellular domain, a single transmembrane domain, and a 277 amino acid cytoplasmic domain. gp130 is a subunit of several heterodimeric cell-surface receptors, including receptors for IL-6, IL-11, IL-27, Oncostatin M, and Leukemia Inhibitory Factor (LIF). The gp130 protein has also been found to exist in a soluble form, which is capable of inhibiting IL-6 signaling. gp130 is expressed mainly on T cells, monocytes, endothelial cells, activated B cells, and plasma cells, and is expressed at lower levels on most leukocytes and epithelial cells. In response to ligand binding, gp130 becomes tyrosine phosphorylated, leading to activation of several signaling pathways including the PI3 kinase, Ras-MAPK and Stat pathways.

Preincubation of mouse splenic T cells with IL-6 is capable of blocking the interaction of KGP130 with gp130, suggesting that the KGP130 binding site lies near the IL-6 interaction domain. Reactivity of KGP130 towards human gp130 has not been observed.

Applications Reported

This KGP130 antibody has been reported for use in flow cytometric analysis.

Applications Tested

This KGP130 antibody has been tested by flow cytometric analysis of mouse splenocytes. This can be used at less than or equal to 0.25 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µl. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

References

Arzt E. gp130 cytokine signaling in the pituitary gland: a paradigm for cytokine-neuro-endocrine pathways. J Clin Invest. 2001 Dec;108(12):1729-33. (PubMed)

Bravo J, Heath JK. Receptor recognition by gp130 cytokines. EMBO J. 2000 Jun 1;19(11):2399-411. (PubMed)

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.ebioscience.com •
info@ebioscience.com

Anti-Mouse CD130 APC

Catalog Number: 17-1302

Also known as: gp130, IL-6 Receptor beta, IL-6RB

RUO: For Research Use Only. Not for use in diagnostic procedures.

Wang XJ, Taga T, Yoshida K, Saito M, Kishimoto T, Kikutani H. gp130, the cytokine common signal-transducer of interleukin-6 cytokine family, is downregulated in T cells in vivo by interleukin-6. Blood. 1998 May 1;91(9):3308-14. (PubMed)

Related Products

11-0031 Anti-Mouse CD3e FITC (145-2C11)

17-4321 Rat IgG2a K Isotype Control APC (eBR2a)