


Anti-Mouse MHC Class II (I-A/I-E) PE-Cyanine7

Catalog Number: 25-5321

Also known as: MHC II, IA, IE, I-A/E, IA/IE

RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of C57Bl/6 splenocytes with Anti-Mouse CD3e APC (cat. 17-0031) and 0.06 ug of Rat IgG2b K Isotype Control PE-Cyanine7 (cat. 25-4031) (left) or 0.06 ug of Anti-Mouse MHC Class II (I-A/I-E) PE-Cyanine7 (right). Cells in the lymphocyte gate were used for analysis.

Product Information

Contents: Anti-Mouse MHC Class II (I-A/I-E)

PE-Cyanine7

Catalog Number: 25-5321 Clone: M5/114.15.2 Concentration: 0.2 mg/mL

Host/Isotype: Rat IgG2b, kappa



Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer **Temperature Limitation:** Store at 2-8°C. Do not

freeze. Light-sensitive material. This tandem dye is sensitive to photo-induced oxidation. Protect this vial from light during storage, handling &



experimental procedures.

Batch Code: Refer to vial

Use By: Refer to vial



Contains sodium azide

Description

The M5/114.15.2 monoclonal antibody reacts with the mouse major histocompatibility complex class II, both I-A and I-E subregion-encoded glycoproteins (I-A b, I-A d, I-A d, I-E d, I-E k, not I-A f, I-A k, or I-A s). It detects a polymorphic determinant present on B cells, monocytes, macrophages, dendritic cells, and activated T lymphocytes from mice carrying the H-2 b, H-2 d, H-2 q, H-2 p, H-2 r and H-2 u but not from mice carrying the H-2 s or H-2 f haplotypes. The M5/114 mAb is reported to inhibit I-A-restricted T cell responses of the H-2 b, H-2 d, H-2 q, H-2 u but not H-2 f, H-2 k, or H-2 s haplotypes.

Applications Reported

This M5/114.15.2 antibody has been reported for use in flow cytometric analysis.

Applications Tested

This M5/114.15.2 antibody has been tested by flow cytometric analysis of mouse splenocytes. This can be used at less than or equal to 0.25 μ g per test. A test is defined as the amount (μ g) of antibody that will stain a cell sample in a final volume of 100 μ L. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

Light sensitivity: This tandem dye is sensitive photo-induced oxidation. Please protect this vial and stained samples from light.

Fixation: Samples can be stored in IC Fixation Buffer (cat. 00-8222) (100 uL cell sample + 100 uL IC Fixation Buffer) or 1-step Fix/Lyse Solution (cat. 00-5333) for up to 3 days in the dark at 4°C with minimal impact on brightness and FRET efficiency/compensation. Some generalizations regarding fluorophore performance after fixation can be made, but clone specific performance should be determined empirically.



Anti-Mouse MHC Class II (I-A/I-E) PE-Cyanine7

Catalog Number: 25-5321

Also known as: MHC II, IA, IE, I-A/E, IA/IE

RUO: For Research Use Only. Not for use in diagnostic procedures.

References

Unternaehrer JJ, Chow A, et al. 2007. The tetraspanin CD9 mediates lateral association of MHC class II molecules on the dendritic cell surface. Proc Natl Acad Sci U S A. 104(1):234-9. (M5/114.15.2, IP, PubMed)

Gueirard, Pascale et al. 2003. "Bordetella bronchiseptica Perists in the Nasal Cavities of Mice and Triggers Early Delivery of Dendritic Cells in the Lymph Nodes Draining the Lower and Upper Respiratory Tract". Infection and Immunity 71(7):4137-4143. (M5/114.15.2, IHC paraffin, PubMed)

Bagavant, Sharp et al. 2002. "Induction and Immunohistology of Autoimmune Ovarian Disease in Cynomolgus Macaques (Macaca fascicularis)". Am J Pathol 160:141-149. (M5/114.15.2, IHC, PubMed)

Li C, Siemasko K, et al. 2002. Cooperative interaction of Ig(alpha) and Ig(beta) of the BCR regulates the kinetics and specificity of antigen targeting. Int Immunol. 14(10):1179-91. (**M5/114.15.2**, IHC frozen and WB, PubMed)

Mediratta SK, Singh N, et al. 1996. Analysis of T-cell hybridomas with an unusual MHC class II-dependent ligand specificity. Immunology. 89(2):238-44. (M5/114.15.2, FA, PubMed)

Germain, R. N., A. Bhattacharya, et al. 1982. A single monoclonal anti-la antibody inhibits antigen-specific T cell proliferation controlled by distinct Ir genes mapping in different H-2 I subregions. J Immunol 128(3): 1409-13.

Bhattacharya, A., M. E. Dorf, et al. 1981. A shared alloantigenic determinant on Ia antigens encoded by the I-A and I-E subregions: evidence for I region gene duplication. J Immunol 127(6): 2488-95.

Related Products

17-0031 Anti-Mouse CD3e APC (145-2C11) 25-4031 Rat IgG2b K Isotype Control PE-Cyanine7

Legal

FOR NON-COMMERCIAL RESEARCH USE ONLY. NOT FOR THERAPEUTIC OR IN VIVO APPLICATIONS. OTHER USE NEEDS LICENSE FROM GE HEALTHCARE BIO-SCIENCES CORP. UNDER U.S. PATENT FOR NON-COMMERCIAL RESEARCH USE ONLY. NOT FOR THERAPEUTIC OR IN VIVO APPLICATIONS. OTHER USE NEEDS LICENSE FROM GE HEALTHCARE BIO-SCIENCES CORP. UNDER U.S. PATENT # 5,268,486, 5,569,587 AND 5,627,027 AND FOREIGN EQUIVALENTS AND PENDING APPLICATIONS. THIS MATERIAL IS SUBJECT TO PROPRIETARY RIGHTS OF GE HEALTHCARE BIO-SCIENCES CORP. AND CARNEGIE MELLON UNIVERSITY AND MADE AND SOLD UNDER LICENSE FROM GE HEALTHCARE BIO-SCIENCES CORP. THIS PRODUCT IS LICENSED FOR SALE ONLY FOR RESEARCH. IT IS NOT LICENSED FOR ANY OTHER USE. THERE IS NO IMPLIED LICENSE HEREUNDER FOR ANY COMMERCIAL USE. COMMERCIAL USE shall include: 1. sale, lease, license or other transfer of the material or any material derived or produced from it; 2. sale, lease, license or other grant of rights to use this Material or any material derived or produced from it; 3. use of this material to perform services for a fee for third parties. IF YOU REQUIRE A COMMERCIAL LICENSE TO USE THIS MATERIAL AND DO NOT HAVE ONE, RETURN THIS MATERIAL, UNOPENED TO EBIOSCIENCE, INC. 10255 SCIENCE CENTER DRIVE, SAN DIEGO, CALIFORNIA 92121 USA AND ANY MONEY PAID FOR THE MATERIAL WILL BE REFUNDED.