

Product Data Sheet

Alexa Fluor® 647 anti-human CD107a (LAMP-1)

Catalog # / Size: 328611 / 25 tests

328612 / 100 tests

Clone: H4A3

Isotype: Mouse IgG1, κ

Workshop Number: P PR-63; BP 473; P P008

Immunogen: Human adult adherent peripheral blood cells

Reactivity: Human, Cross-Reactivity: African Green, Baboon, Chimpanzee, Pigtailed

Macaque, Rhesus

Preparation: The antibody was purified by affinity chromatography, and conjugated with

Alexa Fluor® 647 under optimal conditions. The solution is free of


unconjugated Alexa Fluor® 647.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and

0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from

prolonged exposure to light. Do not freeze.v



Thrombin-activated human peripheral blood platelets were stained with CD107a (clone H4A3) Alexa Fluor® 647 (red histogram) or mouse IgG1, κ Alexa Fluor® 647 (green histogram).

Applications:

Applications: FC - Quality tested

IF - Reported in the literature

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is 5 µl per million cells or 5 µl per 100 µl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Alexa Fluor® 647 has a maximum emission of 668 nm when it is excited at 633nm / 635nm.

** Alexa Fluor® is a registered trademark of Molecular Probes, Inc. Alexa Fluor® dye antibody conjugates are sold under license from Molecular Probes. Inc. for research use only, except for use in combination with microarrays and high content screening, and are covered by pending and issued patents.

Application Notes: Additional reported applications (for the relevant formats) include: Western blotting, immunohistochemical staining², immunofluorescence^{5,7}, and immunoprecipitation⁵.

Application References: 1. Misse D, et al. 1999. Blood 93:2454.

2. Furuta K, et al. 2001. Am. J. Pathol. 159:449. (IHC)

3. Watanabe A, et al. 2011. J. Biol. Chem. 286:10702. PubMed Watariabe A, et al. 2011. 3. Biol. Crieff. 2007. 10702. 1 tabled
Baron Gaillard CL, et al. 2011. Mol. Cell. Biol. 22:5459. PubMed
Hauck CR and Meyer TF. 1997. FEBS Lett. 405:86. (IF, IP)
De Keersmaecker B, et al. 2012. J. Virol. 86:9351. PubMed
Knodler LA, et al. 2010. P. Natl. Acad. Sci. USA. 107:17733. (IF)

Description: CD107a, also known as Lysosome-Associated Membrane Protein 1 (LAMP-1) or LGP-120, is a 110-140 kD type I membrane gly-coprotein. The mature CD107a is heavily glycosylated from a 40 kD core protein. It is expressed by activated platelets, acti-vated lymphocytes, macrophages, epithelial cells, endothelial cells, and some tumor cells. This molecule is located on the luminal side of lysosomes and has been suggested to play a role in the protection of lysosomes membrane from lysosomal hydrolases. Upon activation, CD107a is transferred to the cell membrane surface. It plays a role in cell adhesion and regula-tion of tumor metastasis, and mediates autoimmune disease progression. CD107a is a ligand for galaptin and E-selectin.

Antigen References: 1. Sarafian V, et al. 2006. Arch. Dermatol. Res. 298:7381.

2. Schlossman SF, et al. 1995. Leukocyte Typing V:White Cell Differentiation Antigens. New York:Oxford University

3. Sawada R, et al. 1993. J. Biol. Chem. 268:12675. 4. Chen JW, et al. 1988. J. Biol. Chem. 263:8754. 5. Chen JW, et al. 1986. Biochem. Soc. Symp. 51:97112.

Related Products: Product

Alexa Fluor® 647 Mouse IgG1, κ Isotype Ctrl (FC) MOPC-21 Cell Staining Buffer

RBC Lysis Buffer (10X)

Human TruStain FcX™ (Fc Receptor Blocking Solution)

Clone Application FC, IF FC, ICC, ICFC FC, ICFC FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

