

## **Product Data Sheet**

## APC/Cy7 anti-human HLA-DR

Catalog # / Size: 307617 / 25 tests

307618 / 100 tests

Clone: L243

**Isotype:** Mouse IgG2a, κ

**Reactivity:** Human, **Cross-Reactivity:** African Green, Baboon, Chimpanzee, Common Marmoset, Cotton-topped Tamarin, Cynomolgus, Pigtailed Macaque,

Rhesus, Squirrel Monkey, Dog (Canine)17, 18

**Preparation:** The antibody was purified by affinity chromatography, and conjugated with APC/Cy7 under optimal conditions. The solution is free of unconjugated


APC/Cy7 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and

0.2% (w/v) BSA (origin USA).

Storage: The HLA-DR antibody solution should be stored undiluted at 4°C, and

protected from prolonged exposure to light. **Do not freeze.** 



Human peripheral blood lymphocytes stained with L243 APC/Cy7

## **Applications:**

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. Test size products are transitioning from 20 µl to 5 µl per test. Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 µl staining volume or per 100 µl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Read more at www.biolegend.com/testsize regarding the test size change.

Application Notes: The L243 monoclonal antibody reacts with the HLA-DR antigen, a member of MHC class II molecules. It does not cross react with HLA-DP and HLA-DQ. Additional reported applications (for the relevant formats) include: immunoprecipitation<sup>8</sup>, Western blotting<sup>8</sup>, *in vitro* blocking of mixed lymphocyte reactions<sup>7,9,10</sup>, and immunohistochemical staining of acetone-fixed frozen sections<sup>4,5</sup>. The LEAF™ purified antibody (Endotoxin <0.1 EU/µ g, Azide-Free, 0.2 μm filtered) is recommended for functional assays (Cat. No. 307612). For highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. No. 307648) with a lower endotoxin limit than standard LEAF™ purified antibodies (Endotoxin < 0.01 EU/µg).

> Cy3, Cy5, Cy5.5 and Cy7 are subject to proprietary rights of GE Healthcare Bio-Sciences Corp. and Carnegie Mellon University and made and sold under license from GE Healthcare Bio-Sciences Corp. Sale of this product is licensed for research use only.

- Application References: 1. Brodsky F. 1984. Immunogenetics 19:179.

  - Robbins P, et al. 1987. Human Immunol. 18:301.
     Stites D, et al. 1986. Clin. Immunol. Immunopathol. 38:161.

  - Warnke R, et al. 1980. J. Histochem. Cytochem. 28:771. (IHC)
     Engleman E, et al. 1981. P. Natl. Acad. Sci. USA 78:1791. (IHC)
     Zipf T, et al. 1981. Cancer Res. 41:4786.

7. Goodier M, *et al.* 2000. *J. Immunol.* 165:139. (Block) 8. Esser M, *et al.* 2001. *J. Virol.* 75:6173. (IP, WB) 9. Kalka-Moll WM, *et al.* 2002. *J. Immunol.* 169:6149. (Block)

10. Wang RF, et al. 1999. Science 284:1351. (Block) 11. Zaba LC, et al. 2007. J. Exp. Med. 204:3183. PubMed

12. Fujita H, et al. 2009. P. Natl. Acad. Sci. USA 106:21795. PubMed 13. Charles N, et al. 2010. Nat. Med. 16:701. (FC) PubMed

14. Goncalves RM, et al. 2010. Infect. Immun. 78:4763. PubMed

15. Yoshino N, et al. 2000. Exp. Anim. (Tokyo) 49:97. (FC) 16. Kim WK, et al. 2006. Am. J. Pathol. 168:822. (FC)

17. Stein R, *et al.* 2011. *Leuk. Lymphoma* 52:273. 18. Galkowska H, *et al.* 1996. *Vet. Immunol. Immunopathol.* 53:329.

Description: HLA-DR is a heterodimeric cell surface glycoprotein comprised of a 36 kD α (heavy) chain and a 27 kD β (light) chain.

It is expressed on B cells, activated T cells, monocytes/macrophages, dendritic cells, and other non-professional APCs. In conjunction with the CD3/TCR complex and CD4 molecules, HLA-DR is critical for efficient peptide

presentation to CD4+ T cells.

Antigen References: 1. Levacher M, et al. 1990. Clin. Exp. Immunol. 81:177.

2. Terstappen L, et al. 1990. J. Leukocyte Biol. 48:138.

3. Edwards JA, et al. 1986. J. Immunol. 137:490.



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



van Es A, et al. 1984. Transplantation 37:65.
 O'Doherty U, et al. 1994. Immunology 82:487.
 Thomas R, et al. 1994. J. Immunol. 153:4016.

7. Grouard G, et al. 1996. Nature 384:364.

**Related Products: Product** Clone Application

Cell Staining Buffer
RBC Lysis Buffer (10X)
APC/Cy7 Mouse IgG2a, κ Isotype Ctrl
Human TruStain FcX™ (Fc Receptor **MOPC-173** 

**Blocking Solution**)

FC, ICC, ICFC FC, ICFC FC, ICFC FC, ICC, ICFC



