

## Technical Data Sheet


## FITC Mouse Anti-Human CD3ε

## Product Information

Material Number:	558261
Size:	100 tests
Vol. per Test:	20 µl
Clone:	APA1/1
Isotype:	Mouse IgG1, κ
Reactivity:	QC Testing: Human
Storage Buffer:	Aqueous buffered solution containing BSA and ≤0.09% sodium azide.

## Description

Monoclonal antibody APA1/1 reacts with the intracellular domain of the epsilon chain of the CD3 molecule (CD3-ε). This antibody does not react with the extracellular region of CD3-ε. The assembly of the T cell antigen receptor (TCR)-CD3 complex has been suggested to take place by pairwise interactions of the CD3-ε subunit with either CD3-γ or CD3-δ. These dimers then associate with the TCR heterodimer α/β or γ/δ, and the CD3-ζ homodimer to form the full complex. Studies show that antibodies APA1/1 and SP34 gave a strong reaction with COS cells singly transfected with CD3-ε. Other anti-CD3 antibodies (OKT3, WT31, UCHT1, Leu4) did not react with COS cells singly transfected with CD3-ε. This reagent could be useful for the study of T cell development or the study of conformational changes of CD3-ε upon ligand binding to TCR-CD3 complex.



*CD3-epsilon, clone APA1/1, intracellular reactivity on permeabilized peripheral blood lymphocytes fixed and permeabilized with 70-80% cold ethanol and then analyzed by flow cytometry. Recommended isotype control: FITC mouse IgG1, κ mAb MOPC-21, Cat. No. 555909.*

## Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with FITC under optimum conditions, and unreacted FITC was removed.

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

## Application Notes

## Application

Intracellular staining (flow cytometry)	Routinely Tested
---	------------------

## Recommended Assay Procedure:

## Protocol for PBMC cells fixation, permeabilization and staining

1. Harvest, count and pellet PBMC cells following standard procedures.
2. While vortexing, add 5 ml cold 70% - 80% ethanol dropwise into the cell pellet ( $1-5 \times 10^7$  cells). Incubate at -20°C for at least 2 hours. These fixed cells can be stored at -20°C for up to 60 days prior to staining.
3. Wash twice with 30-40 ml staining buffer (PBS with 1% FBS, 0.09% NaN<sub>3</sub>), centrifuge for 10 minutes at 200 x g.
4. Resuspend the cells to a concentration of  $1 \times 10^6$ /ml.
5. Transfer 100 µl ( $1 \times 10^6$  cells) cell suspension into each sample tube.

## BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit [bdbiosciences.com/how\\_to\\_order/](http://bdbiosciences.com/how_to_order/)

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2008 BD



6. Add 20 µl of properly diluted fluorescence conjugated antibody into the tubes above. Mix gently.
7. Incubate the tubes at room temperature (RT) for 20-30 minutes in the dark.
8. Wash with 2 ml of staining buffer at 200 x g for 5 minutes.
9. Aspirate the supernatant.
10. Add 0.5 ml of staining buffer to each tube.
11. Proceed to flow cytometric analysis.

### Suggested Companion Products

Catalog Number	Name	Size	Clone
555909	FITC Mouse IgG1, κ Isotype Control	100 tests	MOPC-21

### Product Notices

1. This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use  $1 \times 10^6$  cells in a 100-µl experimental sample (a test).
2. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
3. Please refer to [www.bdbiosciences.com/pharming/en/protocols](http://www.bdbiosciences.com/pharming/en/protocols) for technical protocols.
4. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
5. Source of all serum proteins is from USDA inspected abattoirs located in the United States.

### References

Barroto A, Mallabiarrena A, Albar JP, Martinez-A C, Alarcon B. Characterization of the region involved in CD3 pairwise interactions within the T cell receptor complex. *J Biol Chem.* 1998; 273:12807-12816.(Biology)

Gil D, Schamel WWA, Montoya M, Sanchez-Madrid F, and Alarcon B. Recruitment of Nck by CD3-epsilon reveals a ligand-induced conformational change essential for T cell receptor signaling and synapse formation. *Cell.* 2002; 109:901-912.(Biology)

Slamerson A, Sanchez-Madrid F, Ursa MA, Fresno M, and Alarcon B. A conformational epitope expressed upon association of the CD3-epsilon with either CD3-delta or CD3-gamma is the main target for recognition by anti-CD3 monoclonal antibodies. *J Immunol.* 1991; 147:3047-3052.(Biology)