


Technical Data Sheet

Biotin Hamster Anti-Mouse CD27

Product Information	
Material Number:	558753
Size:	0.1 mg
Concentration:	0.5 mg/ml
Clone:	LG.3A10
Immunogen:	Armenian hamster fibroblast line ARHO12 transfected with mouse Cd27 cDNA
Isotype:	Armenian Hamster IgG1, κ
Reactivity:	QC Testing: Mouse Tested in Development: Rat
Storage Buffer:	Aqueous buffered solution containing ≤0.09% sodium azide.

Description

The LG.3A10 antibody reacts with CD27, a lymphocyte-restricted member of the Tumor Necrosis Factor Receptor family which binds to CD70. The CD27 molecule is a 45-kDa transmembrane glycoprotein which is constitutively expressed by lymphocytes of the T lineage: virtually all thymocytes and over 90% of peripheral T cells bearing both αβ and γδ T-cell receptors. CD27 cooperates with the pre-TCR in mediating thymocyte differentiation and expansion. In addition, one to ten percent of mature peripheral B cells express CD27, and CD27's role in the differentiation of human plasma cells has been studied. Mouse NK cells, freshly isolated and IL-2-activated, also express CD27. In the bone marrow, CD27 is found on a progenitor population which provides short-term hematopoietic reconstitution. Cells of the myeloid lineage do not express CD27. Cross-linked LG.3A10 mAb has been reported to amplify the proliferative response of purified T lymphocytes to suboptimal mitogenic stimulation¹ and to enhance NK-cell proliferation and IFN-γ production. In contrast, non-cross-linked LG.3A10 mAb inhibits CD3-induced pre-T cell development by interfering with the receptor-ligand interaction. This hamster mAb to a mouse leukocyte antigen has been observed to cross-react with a similar population of rat leukocytes.



Expression of CD27 on mouse splenocytes. C57BL/6 splenocytes were stained simultaneously with PE-conjugated anti-mouse CD3e mAb 500A2 (Cat. No. 553240) and biotinylated mAb LG.3A10 (right panel), followed by Avidin-FITC (Cat. No. 554057). Flow cytometry was performed on a BD FACScan™ flow cytometry system.

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. The antibody was conjugated with biotin under optimum conditions, and unreacted biotin was removed. Store undiluted at 4° C and protected from prolonged exposure to light. Do not freeze.

Application Notes

Application	
Flow cytometry	Routinely Tested

BD Biosciences

www.bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit www.bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.
For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.
BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2007 BD



Suggested Companion Products

Catalog Number	Name	Size	Clone
553240	PE Hamster Anti-Mouse CD3e	0.1 mg	500A2
554057	Avidin FITC	0.5 mg	(none)
553970	Biotin Hamster IgG1 κ Isotype Control	0.25 mg	A19-3

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
4. Although hamster immunoglobulin isotypes have not been well defined, BD Biosciences Pharmingen has grouped Armenian and Syrian hamster IgG monoclonal antibodies according to their reactivity with a panel of mouse anti-hamster IgG mAbs. A table of the hamster IgG groups, Reactivity of Mouse Anti-Hamster Ig mAbs, may be viewed at http://www.bdbiosciences.com/pharmingen/hamster_chart_11x17.pdf.

References

Agematsu K, Hokibara S, Nagumo H, Shinozaki K, Yamada S, Komiyama A. Plasma cell generation from B-lymphocytes via CD27/CD70 interaction. *Leuk Lymphoma*. 1999; 35(3-4):219-225.(Biology)

Gravestien LA, Blom B, Noltén LA, et al. Cloning and expression of murine CD27: comparison with 4-1BB, another lymphocyte-specific member of the nerve growth factor receptor family. *Eur J Immunol*. 1993; 23(4):943-950.(Biology)

Gravestien LA, Nieland JD, Kruisbeek AM, Borst J. Novel mAbs reveal potent co-stimulatory activity of murine CD27. *Int Immunol*. 1995; 7(4):551-557. (Immunogen)

Gravestien LA, van Ewijk W, Ossendorp F, Borst J. CD27 cooperates with the pre-T cell receptor in the regulation of murine T cell development. *J Exp Med*. 1996; 184(2):675-685.(Biology)

Takeda K, Oshima H, Hayakawa Y, et al. CD27-mediated activation of murine NK cells. *J Immunol*. 2000; 164(4):1741-1745.(Biology)

Wiesmann A, Phillips RL, Mojica M, et al. Expression of CD27 on murine hematopoietic stem and progenitor cells. *Immunity*. 2000; 12(2):193-199.(Biology)