Technical Data Sheet

BV421 Rat Anti-Mouse Siglec-F

Product Information

Material Number: 562681

Alternate Name: Siglec5; sialic acid binding Ig-like lectin 5; SIGL5; Siglec-5; CD170

 Size:
 50 µg

 Concentration:
 0.2 mg/ml

 Clone:
 E50-2440

Immunogen: Mouse Siglec-F and human IgG Fc recombinant fusion protein


 $\begin{array}{lll} \textbf{Isotype:} & \text{Rat (LOU) IgG2a, } \kappa \\ \textbf{Reactivity:} & \text{QC Testing: Mouse} \end{array}$


Storage Buffer: Aqueous buffered solution containing BSA and ≤0.09% sodium azide.

Description

The E50-2440 antibody reacts with Siglec-F, the sixth siglec protein to be reported in the mouse. Siglecs are the sialic acid-binding immunoglobulin superfamily <u>lec</u>tins defined in the human, each of which has a distinctive expression pattern in the hematopoietic system and at least some of which are known to mediate cell-cell interactions. Orthologous proteins of human Siglec-1 (Sialoadhesin or CD169), Siglec-2 (CD22), and Siglec-4 (myelin-associated glycoprotein) have been characterized in the mouse. Human Siglec-3 (CD33) and Siglecs-5 through -10 are encoded by a cluster of closely related genes, and each has two cytoplasmic ITIM (Immunoreceptor Tyrosine-based Inhibitory Motifs). Similarly, mouse Siglec-F is encoded by a gene in a syntenic cluster in the mouse, and the protein has sialic acid-binding activity and an intracytoplasmic ITIM. Its expression pattern differs from those of the human Siglec-3-related proteins in that it is found on immature cells of the myelomonocytic lineage, with reduced expression on mature neutrophils and monocytes, and not on lymphoid cells. It has been proposed that mAb E50-2440 may be used for identification of immature myelomonocytic cells in the mouse.

The antibody was conjugated to BD HorizonTM BV421 which is part of the BD HorizonTM Brilliant VioletTM family of dyes. With an Ex Max of 407-nm and Em Max at 421-nm, BD HorizonTM BV421 can be excited by the violet laser and detected in the standard Pacific BlueTM filter set (eg, 450/50-nm filter). BD HorizonTM BV421 conjugates are very bright, often exhibiting a 10 fold improvement in brightness compared to Pacific BlueTM conjugates.





Multicolor flow cytometric analysis of Siglec-F expression on BALB/c mouse bone marrow cells. Bone marrow cells were stained simultaneously with FITC Rat Anti-Mouse CD11b antibody (Cat. No. 553310/557396/561688) and with either BD Horizon™ BV421 Rat IgG2a, κ Isotype Control (Cat. No. 562602; Left Panel) or BD Horizon™ BV421 Rat Anti-Mouse Siglec-F (Cat. No. 562681; Right Panel) in the presence of Purified Rat Anti-Mouse CD16/CD32 antibody (Mouse BD Fc Block™) (Cat. No. 553141/553142). Two-color flow cytometric dot plots show the correlated expression patterns of Siglec-F (or Ig isotype control staining) versus CD11b for gated events with the forward and side light-scatter characteristics of viable bone marrow cells. Flow cytometry was performed using a BD™ LSR II Flow Cytometer System.

Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with BD Horizon™ BV421 under optimum conditions, and unconjugated antibody and free BD Horizon™ BV421 were removed.

BD Biosciences

bdbiosciences.com

 United States
 Canada
 Europe
 Japan
 Asia Pacific
 Latin America/Caribbean

 877.232.8995
 800.979.9408
 32.53.720.550
 0120.8555.90
 65.6861.0633
 55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be constructed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be help responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is stictly prohibited. For Research Use Only, Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD



Application Notes

Application

Flow cytometry	Routinely Tested

Suggested Companion Products

Catalog Number	<u>Name</u>	Size	Clone	
554656	Stain Buffer (FBS)	500 ml	(none)	
562602	BV421 Rat IgG2a, κ Isotype Control	50 μg	R35-95	
553141	Purified Rat Anti-Mouse CD16/CD32 (Mouse BD Fc Block™)	0.1 mg	2.4G2	
553142	Purified Rat Anti-Mouse CD16/CD32 (Mouse BD Fc Block™)	0.5 mg	2.4G2	
553310	FITC Rat Anti-Mouse CD11b	0.5 mg	M1/70	
557396	FITC Rat Anti-Mouse CD11b	0.1 mg	M1/70	
561688	FITC Rat Anti-Mouse CD11b	25 μg	M1/70	

Product Notices

- Since applications vary, each investigator should titrate the reagent to obtain optimal results.
- 2. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
- 3. An isotype control should be used at the same concentration as the antibody of interest.
- 4. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
- Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
- For fluorochrome spectra and suitable instrument settings, please refer to our Multicolor Flow Cytometry web page at www.bdbiosciences.com/colors.
- 7. Pacific BlueTM is a trademark of Molecular Probes, Inc., Eugene, OR.
- 8. Brilliant VioletTM 421 is a trademark of Sirigen.

References

Angata T, Hingorani R, Varki NM, Varki A. Cloning and characterization of a novel mouse Siglec, mSiglec-F: differential evolution of the mouse and human (CD33) Siglec-3-related gene clusters. *J Biol Chem.* 2001; 276(48):45128-45136. (Immunogen: Flow cytometry, Fluorescence activated cell sorting) Crocker PR, Varki A. Siglecs, sialic acids and innate immunity. *Trends Immunol.* 2001; 22(6):337-342. (Biology)

BD Biosciences

bdbiosciences.com

 United States
 Canada
 Europe
 Japan
 Asia Pacific
 Latin America/Caribbean

 877.232.8995
 800.979.9408
 32.53.720.550
 0120.8555.90
 65.6861.0633
 55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be constructed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be help responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is stictly prohibited. For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD



562681 Rev. 2 Page 2 of 2