

## Technical Data Sheet

## BV510 Rat Anti-Pax-5


## Product Information

<b>Material Number:</b>	<b>563191</b>
<b>Alternate Name:</b>	Pax5, PAX5, KLP, BSAP, EBB-1; B-cell-specific transcription factor
<b>Size:</b>	50 tests
<b>Vol. per Test:</b>	5 µl
<b>Clone:</b>	1H9
<b>Immunogen:</b>	Recombinant Mouse Pax-5 protein containing aa 154-284
<b>Isotype:</b>	Rat IgG2a, κ
<b>Reactivity:</b>	QC Testing: Mouse Tested in Development: Human
<b>Storage Buffer:</b>	Aqueous buffered solution containing BSA and ≤0.09% sodium azide.

## Description

The 1H9 monoclonal antibody clone specifically binds to human and mouse Paired box protein Pax-5. Pax-5 is a ~50 kDa protein that is also known as B-cell-specific transcription factor and B cell specific activator protein (BSAP). Pax-5 is a member of the paired box (Pax) family of transcription factors. It is expressed in pro-B, pre-B and mature B cells. Through the Groucho family of co-repressors, Pax-5 likely functions as a transcriptional repressor of non-B-lymphoid genes during the B cell commitment process. In the early stages of B cell development, Pax-5 influences the expression of several B-cell-specific genes, such as CD19 and CD20 and maintains B cell identity. Pax-5 suppression is involved in the upregulation of Blimp-1 leading to the development of Pax-5-negative plasma cells. Pax-5 mRNA is transiently detected in the mesencephalon and spinal cord during embryogenesis. Expression then shifts to the fetal liver and correlates with the onset of B lymphopoiesis. Altered forms and expression patterns of Pax-5 have been associated with some lymphoid and nonlymphoid cancers.

The antibody was conjugated to BD Horizon™ BV510 which is part of the BD Horizon™ Brilliant Violet™ family of dyes. With an Ex Max of 405-nm and Em Max at 510-nm, BD Horizon™ BV510 can be excited by the violet laser and detected in the BD Horizon™ V500 (525/50-nm) filter set. BD Horizon™ BV510 conjugates are useful for the detection of dim markers off the violet laser.



**Multicolor flow cytometric analysis of Pax5 expression in human and mouse leucocytes.** Human peripheral blood mononuclear cells (PBMC) and BALB/c mouse splenic leucocytes were fixed and permeabilized using the BD Pharmingen™ Transcription Factor Buffer Set (Cat. No. 562574/562725). The cells were then stained with either BD Horizon™ BV510 Rat Anti-Mouse Pax5 antibody (Cat. No. 563191) or BD Horizon™ BV510 Rat IgG2a, κ Isotype Control (Cat. No. 562952). The PBMC were counterstained with PE Mouse Anti-Human CD21 antibody (Cat. No. 555422/557327/561768). The mouse splenic leucocytes were counterstained with APC Rat Anti-Mouse CD45R/B220 antibody (Cat. No. 553092/561880). Flow cytometric analysis was performed using a BD LSR™ II Flow Cytometer System.

**Panel A. Human Peripheral Blood Lymphocytes:** The two-color flow cytometric dot plots show the correlated expression patterns of Ig Isotype control staining (Left Plot) and Pax5 (Right Plot) versus CD21 for events with the forward and side light-scatter characteristics of intact human peripheral blood lymphocytes.

**Panel B. Mouse Splenocytes:** The dot plots show the coexpression patterns of Ig Isotype control staining (Left Plot) or Pax5 (Right Plot) versus CD45R/B220 for events with the light-scatter characteristics of intact mouse splenocytes.

## Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with BD Horizon™ BV510 under optimum conditions, and unconjugated antibody and free BD Horizon™ BV510 were removed.

## BD Biosciences

[bdbiosciences.com](http://bdbiosciences.com)

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	800.979.9408	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country contact information, visit [bdbiosciences.com/contact](http://bdbiosciences.com/contact)

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD



## Application Notes

### Application

Intracellular staining (flow cytometry)

Routinely Tested

### Suggested Companion Products

Catalog Number	Name	Size	Clone
554656	Stain Buffer (FBS)	500 ml	(none)
562952	BV510 Rat IgG2a, $\kappa$ Isotype Control	50 $\mu$ g	R35-95
555899	Lysing Buffer	100 ml	(none)
562574	Transcription Factor Buffer Set	100 tests	(none)
562725	Transcription Factor Buffer Set	25 tests	(none)
555422	PE Mouse Anti-Human CD21	100 tests	B-ly4
557327	PE Mouse Anti-Human CD21	50 tests	B-ly4
561768	PE Mouse Anti-Human CD21	25 tests	B-ly4
553092	APC Rat Anti-Mouse CD45R/B220	0.1 mg	RA3-6B2
561880	APC Rat Anti-Mouse CD45R/B220	25 $\mu$ g	RA3-6B2

### Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
3. An isotype control should be used at the same concentration as the antibody of interest.
4. Please refer to [www.bdbiosciences.com/pharming/protocols](http://www.bdbiosciences.com/pharming/protocols) for technical protocols.
5. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
6. For fluorochrome spectra and suitable instrument settings, please refer to our Multicolor Flow Cytometry web page at [www.bdbiosciences.com/colors](http://www.bdbiosciences.com/colors).
7. Brilliant Violet™ 510 is a trademark of Sirigen.

### References

Adams B, Dorfler P, Aguzzi A, et al. Pax-5 encodes the transcription factor BSAP and is expressed in B lymphocytes, the developing CNS, and adult testis. *Genes Dev.* 1992; 6(9):1589-1607. (Biology)

Kallies A, Hasbold J, Fairfax K, et al. Initiation of plasma-cell differentiation is independent of the transcription factor Blimp-1. *Immunity.* 2007; 26(5):555-566. (Immunogen: Western blot)

Klein U, Tu Y, Stolzovitzky GA, et al. Transcriptional analysis of the B cell germinal center reaction. *Proc Natl Acad Sci U S A.* 2003; 100(5):2639-2644. (Biology)

O'Brien P, Morin P, Jr., Ouellette RJ, Robichaud GA. The Pax-5 gene: a pluripotent regulator of B-cell differentiation and cancer disease. *Cancer Res.* 2011; 71(24):7345-7350. (Biology)

Zwollo P, Arrieta H, Ede K, Molinder K, Desiderio S, Pollock R. The Pax-5 gene is alternatively spliced during B-cell development. *J Biol Chem.* 1997; 272(15):10160-10168. (Biology)

## BD Biosciences

[bdbiosciences.com](http://bdbiosciences.com)

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	800.979.9408	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country contact information, visit [bdbiosciences.com/contact](http://bdbiosciences.com/contact)

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD

