# **Technical Data Sheet**

# **Purified Mouse Anti-XIAP**

#### **Product Information**

 Material Number:
 610762

 Alternate Name:
 hILP

 Size:
 50 μg

 Concentration:
 250 μg/ml

 Clone:
 48/hILP/XIAP

Immunogen: Human hILP aa. 268-426

Isotype: Mouse IgG1

Reactivity: QC Testing: Human

Tested in Development: Rat, Mouse, Frog

Target MW: 57 kD


Storage Buffer: Aqueous buffered solution containing BSA, glycerol, and ≤0.09% sodium


azide.

### Description

Apoptosis is a genetically programmed, selective process of cell death that occurs during normal cell differentiation and development of multicellular organisms. Viruses depend on the biosynthetic machinery of their host cell for the production of progeny and survival. Therefore, many viruses encode proteins that protect the cell from apoptosis. hILP (human IAP-like protein) is a human homologue of the viral IAP (Inhibitor of Apoptosis Protein). hILP is a widely expressed cytoplasmic protein of 497 amino acids with three BIR (Baculovirus IAP repeats) domains and a C-terminal RING finger domain. hILP-transfected cells are protected against the apoptotic effects of Sindbis virus infection and ICE (interleukin-1β converting enzyme) expression. This product is sold under license from Aegera Therapeutics, Inc.

This antibody is routinely tested by western blot analysis. Other applications were tested at BD Biosciences Pharmingen during antibody development only or reported in the literature.





Western blot analysis of hILP/XIAP on Jurkat lysate. Lane 1: 1:4000, lane 2: 1:8000, lane 3: 1:12000 dilution of anti-XIAP. Immunofluorescent staining of Rabbit Liver tissue with anti-XIAP

### **Preparation and Storage**

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. Store undiluted at  $-20^{\circ}$  C.

## **BD Biosciences**

www.bdbiosciences.com

United States Canada Europe Japan Asia Pacific Latin America/Caribbean 877.232.8995 888.259.0187 32.53.720.550 0120.8555.90 65.6861.0633 55.11.5185.9995 For country-specific contact information, visit www.bdbiosciences.com/how\_to\_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only, Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2007 BD

**BD** 

610762 Rev. 1 Page 1 of 2

### **Application Notes**

### Application

Western blot	Routinely Tested
Immunofluorescence	Tested During Development
Immunohistochemistry	Tested During Development
Immunoprecipitation	Not Recommended

### **Suggested Companion Products**

Catalog Number	Name	Size	Clone	
611451	Jurkat Cell Lysate	500 μg	(none)	_
554001	FITC Goat Anti-Mouse Igs (Multiple Adsorption)	0.5 mg	Polyclonal	
554002	HRP Goat Anti-Mouse Igs	1.0 ml	(none)	

### **Product Notices**

- Since applications vary, each investigator should titrate the reagent to obtain optimal results.
- Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
- 3. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
- Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

#### References

Deng Y, Lin Y, Wu X. TRAIL-induced apoptosis requires Bax-dependent mitochondrial release of Smac/DIABLO. Genes Dev. 2002; 16(1):33-45.(Clone-specific: Immunoprecipitation, Western blot)

Duckett CS, Nava VE, Gedrich RW. A conserved family of cellular genes related to the baculovirus iap gene and encoding apoptosis inhibitors. EMBO J. 1996; 15(11):2685-2694.(Biology)

Fulda S, Meyer E, Debatin KM. Metabolic inhibitors sensitize for CD95 (APO-1/Fas)-induced apoptosis by down-regulating Fas-associated death domain-like interleukin 1-converting enzyme inhibitory protein expression. Cancer Res. 2000; 60(14):3947-3956.(Clone-specific: Western blot)

Martins LM, laccarino I, Tenev T. The serine protease Omi/HtrA2 regulates apoptosis by binding XIAP through a reaper-like motif. J Biol Chem. 2002; 277(1):439-444.(Clone-specific: Western blot)

Tamm I, Kornblau SM, Segall H. Expression and prognostic significance of IAP-family genes in human cancers and myeloid leukemias. Clin Cancer Res. 2000; 6(5):1796-1803.(Clone-specific: Western blot)

## **BD Biosciences**

www.bdbiosciences.com

United States Canada **Asia Pacific** Latin America/Caribbean Europe Japan 877.232.8995 888.259.0187 32.53.720.550 0120.8555.90 65.6861.0633 55.11.5185.9995

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2007 BD



